

SPECTRAL MULTIPLIERS FOR SCHRÖDINGER OPERATORS WITH PÖSCHL-TELLER POTENTIAL

SHIJUN ZHENG

ABSTRACT. We prove a sharp Mihlin-Hörmander multiplier theorem for Schrödinger operators H on $\mathbb{R}^n$. The method, which allows us to deal with general potentials, improves Hebisch's method relying on heat kernel estimates for positive potentials [22, 12]. Our result applies to, in particular, the *negative* Pöschl-Teller potential $V(x) = -\nu(\nu + 1) \operatorname{sech}^2 x$, $\nu \in \mathbb{N}$, for which H has a resonance at zero.

1. INTRODUCTION

Spectral multiplier theorem for differential operators plays a significant role in harmonic analysis and PDEs. It is closely related to the study of the associated function spaces and Littlewood-Paley theory. Let $H = -\Delta + V$ be a Schrödinger operator on $\mathbb{R}^n$, where V is real-valued. Spectral multipliers for H have been considered in [22, 16, 14, 15, 3] and [12] for positive potentials. The case of negative potential is quite different and is not covered by the methods in these papers. Resonance and eigenvalue can occur that makes the analysis more involved. In this paper we are mainly concerned with proving a Mihlin-Hörmander type multiplier theorem on L^p spaces for the Schrödinger operator with the negative P-T potential

$$(1) \quad V(x) = -\nu(\nu + 1) \operatorname{sech}^2 x, \quad \nu \in \mathbb{N}.$$

In [31, 47] we are able to extend the sharp spectral multiplier theorem on Triebel-Lizorkin spaces by modifying the argument in this note.

Spectral multiplier problem requires both high and low energy analysis. In high energy the kernel of the multiplier operator $m(H)$ can be controlled by a weighted L^2 estimates. In low energy, roughly, it can be controlled pointwise by an approximation to the identity. This is

Date: September 29, 2006.

2000 Mathematics Subject Classification. Primary: 42B25; Secondary: 35J10, 35P25, 35Q40.

Key words and phrases. spectral multiplier, Schrödinger operator, Littlewood-Paley theory.

The author is supported by DARPA grant HM1582-05-2-0001.

the idea when dealing with positive selfadjoint operators [22, 12] where there is a rough kernel.

However for negative operator in the Schrödinger case, resonance and eigenvalue can occur even for smooth and rapidly decaying potentials, which lead to failure of the pointwise control of the kernel in lower energy. The purpose of this paper is to develop a general treatment to overcome this difficulty. We find that this pointwise estimate can be substituted with a weaker estimate (in integral form) that turns out still work. This is the approach we will apply to the Pöschl-Teller potential model. The P-T potential arises in standing wave problem for the cubic wave equation

$$(2) \quad -u_{tt} + u_{xx} + 2u - u^3 = 0.$$

In [47] we give a general treatment on spectral multiplier problem for Schrödinger operators satisfying for every $j \in \mathbb{Z}$

$$(3) \quad |\Phi_j(H)(x, y)| \leq c_n \frac{2^{nj/2}}{(1 + 2^{j/2}|x - y|)^{n+\epsilon}}.$$

where $\Phi_j(x) = \Phi(2^{-j}x)$, $\Phi \in C_0^\infty(\mathbb{R})$.

The assumption is verified when H is a Schrödinger operator $-\Delta + V$, $V \geq 0$ is in $L_{loc}^1(\mathbb{R}^n)$ [32, 22] or H is a uniformly elliptic operator on $L^2(\mathbb{R}^n)$ [9, Theorem 3.4.10]. It is showed in [22, 46] that the decay (3) is satisfied whenever the heat kernel of e^{-tH} satisfies the upper Gaussian bound

$$0 \leq e^{-tH}(x, y) \leq c_n t^{-n/2} e^{-c|x-y|^2/t}.$$

However, when V is negative, eigenvalue and resonance may occur at the origin. the seemingly ubiquitous decay (3) for general selfadjoint operators is not valid for all j . Our approach shows that if (3) replaced by an integral version (1), the argument in [22, 12] can still work. This treatment will be further elaborated in the study of spectral calculus for rough potentials in the critical class in a following paper.

The basic ingredients we need to show are two weighted inequalities

- (a) If $\Phi \in C_0^\infty(\mathbb{R})$ there exists a finite measure μ such that for each interval I with length $2^{-j/2}$, $j \in \mathbb{Z}$,

$$\begin{aligned} |\Phi_j(H)(x, y)| &\leq c \int_{u \in \mathbb{R}^n} \frac{2^{jn/2}}{(1 + 2^{j/2}|x - y - u|)^{n+\epsilon}} d\mu(u) \\ &\approx \rho_j * \mu(x - y) \end{aligned}$$

where $\rho_j(x) := 2^{jn/2}(1 + 2^{j/2}|x|)^{-n-\epsilon}$.

(b)

$$\sup_y \|\langle 2^j(x-y) \rangle^\alpha \phi_j(H)(x, y)\|_{L_x^2} \leq C \sup_j \|\chi \phi(2^{-j}\xi^2)\|_{X^\alpha}$$

where X^α is $C^\alpha(\mathbb{R})$ or W^α , $\alpha > n/2$. χ is a given smooth cut-off function with support off 0.

Observe that (a) is a weaker condition than the pointwise estimate for the decay of $\phi_j(H)$. Indeed, the pointwise decay (3) is a special case when $\mu = \delta$.

1.1. Weighted L^2 estimates for kernel of $m(H)$. There are a few ways to prove L^p boundedness for $m(H)$.

The usual condition is the Hörmander integral condition

$$\int_{|x-y|>2|x-\bar{y}|} |m(H)(x, y) - m(H)(x, \bar{y})| dx \leq A$$

for $y \in I$, $\bar{y}$ the center of I , I being any cube in $\mathbb{R}^n$. This is what is shown in [3]. For P-T potential similar estimate is not valid in low energy. This is the reason why we need to consider the weighted L^2 estimate in (b).

The other way is to use wave operator method [41, 42]. However wave operator method does not give the sharper weak (1, 1) result.

2. MAIN RESULTS

Let H be a selfadjoint operator on $L^2(\mathbb{R}^n)$. Then if $\phi \in L^\infty$, we can define $\phi(H) = \int \phi(\lambda) dE_\lambda$ by functional calculus, where $H = \int \lambda dE_\lambda$ is the spectral resolution of H .

Our main result is the following

Theorem 2.1. *Suppose H verifies the weighted decay (a) and weighted L^2 inequality (b). Then $m(H)$ is bounded on $L^p(\mathbb{R}^n)$, $1 < p < \infty$ and of weak type (1, 1). Moreover,*

$$\|m(H)\|_{L^1 \rightarrow w-L^1} \leq C(m)$$

$$C(m) := \|m\|_\infty + \sup_{\lambda > 0} \|\chi(\cdot)m(\lambda \cdot)\|_{X^\alpha}.$$

Remark. One may view condition (a) as a “pointwise” control of the kernel in *lower* energy while condition (b) as a norm control in *higher* energy.

The conditions of Theorem 2.1 applies to the case when the kernel of $\phi(H)$ is slowly decaying or lack of smoothness. It can also simplifies the proof of some known results on other potential of polynomial growth, e.g. the Hermite operator on $\mathbb{R}^n$ [47].

Applying the theorem to P-T potential we obtain

Theorem 2.2. *Let $H = -d^2/dx^2 + V_\nu$, $\nu \in \mathbb{N}$, where V_ν is the P-T potential in (1). Then H satisfies the weighted decay (a) and weighted L^2 inequality (b) with $X^\alpha = C^\alpha$, $\alpha = 1$. Therefore the conclusion of Theorem 2.1 holds for the one dimensional P-T model.*

From section 5 we know the derivative of the kernel fail to satisfy nice decay, making it difficult to control the difference of $m(H)$ and thus leading to the failing of low energy estimates for the Hörmander integral condition.

A recently developed approach [12] by Sikora et al extended Hebisch's method [22] that apply to positive operators efficiently but rely heavily on heat kernel estimates. However, in dimensions one and two when the potential is negative, such a heat kernel estimate is NOT available. Therefore we consider more direct approach and would rather state and prove the multiplier result in Theorem 2.1 for general dimensions.

We will assume $\Phi, \varphi \in C_0^\infty(\mathbb{R})$ satisfy the condition

$$(4) \quad \sum_{j=-\infty}^{\infty} \varphi_j(x) = 1 \quad x \neq 0$$

$$(5) \quad \Phi(x) + \sum_{j=1}^{\infty} \varphi_j(x) = 1, \quad \forall x$$

where $\varphi_j(x) = \varphi(2^{-j}x)$.

3. PROOF OF THEOREM 2.1

The technical lemmas we need in proving Theorem 2.1 are:

Lemma 3.1. *Let $y \in I$, $I \subset \mathbb{R}^n$ a cube with length $t = \ell(I)$. Let $2^{-jI/2} \sim t$. Then (a) For $t > 0, j \in \mathbb{Z}$,*

$$\int_{|x-y| \geq 2t} |m_j(H)(1 - \Phi_{j_I}(H))(x, y)| dx \leq C(2^{j/2}t)^{\frac{n}{2}-s} \|m(\xi^2)\|_{\dot{H}^s}$$

s > n/2. (b)

$$\int_{|x-y| \geq 2t} \sum_{j=-\infty}^{\infty} |m_j(H)(1 - \Phi_{j_I}(H))(x, y)| dx \leq A.$$

In particular,

$$\int_{|x-y| \geq 2t} \sup_{j \in \mathbb{Z}} |m_j(H)(1 - \Phi_{j_I}(H))(x, y)| dx \leq A.$$

Lemma 3.2. *Condition (b) implies*

$$\begin{aligned} \max_{y \in I} |\Phi_j(H)(x, y)| &\leq c \min_{y \in I} \int_{u \in \mathbb{R}^n} \frac{2^{jn/2}}{(1 + 2^{j/2}|x - y - u|)^{n+\epsilon}} d\mu(u) \\ &\approx \min_{y \in I} \rho_j * \mu(x - y) \\ &\leq \frac{1}{|I|} \int_{z \in I} \rho_j * \mu(x - z) dz \end{aligned}$$

where $\rho_j(x) := 2^{jn/2}(1 + 2^{j/2}|x|)^{-n-\epsilon}$.

The proof is based on the observation if $\ell(I) = r_I$, the side length of a cube I in $\mathbb{R}^n$,

$$\sup_{y \in I} (1 + |x - y|)^{-n-\epsilon} \leq C \min_{y \in I} (1 + |x - y|)^{-n-\epsilon}$$

hence

$$\sup_{y \in I} (1 + |x - y|/t)^{-n-\epsilon} \leq C \min_{y \in I} (1 + |x - y|/t)^{-n-\epsilon} \leq \frac{C}{|I|} \int_I (1 + |x - y|/t)^{-n-\epsilon} dy$$

where $t \sim \ell(I)$, I is any cube, see [22].

Proof of weak (1,1). If applying the C-Z decomposition we know the main part is how to handle the “bad” function $b = \sum_k b_k$ where $b_k \subset I_k$, I_k being disjoint intervals in $\mathbb{R}$.

Proof. Let $\Phi \subset [-1, 1]$, $\Phi_j(x) = \Phi(2^{-j}x)$. Write

$$m(H)b(x) = \sum_k m(H)(1 - \Phi_{j_k}(H))b_k(x) + \sum_k m(H)\Phi_{j_k}(H)b_k(x).$$

where $2^{-j_k} \sim \ell(I_k)^2$. We need to show

$$\begin{aligned} &|\{x \in \mathbb{R} \setminus \cup_k I_k^* : |m(H)b(x)| > \lambda/2\}| \\ &\leq |\{x \in \mathbb{R} \setminus I_k^* : \sum_k |m(H)(1 - \Phi_{j_k}(H))b_k(x)| > \lambda/4\}| \\ &\quad + |\{x \in \mathbb{R} \setminus I_k^* : \sum_k |m(H)\Phi_{j_k}(H)b_k(x)| > \lambda/4\}| \\ &\leq \lambda^{-1} \|f\|_1, \end{aligned}$$

where $b = \sum_k b_k$ (convergence in $L^1 \cap L^q$ so $Tb(x) = \sum_k Tb_k$ in L^q)

Higher energy

Denote I_k^* the cube having length three times the length of I_k with the same center as I_k . If $x \notin \cup_k I_k^*$, $I_k \subset \{y : |y - x| > r_k\}$, r_k being

the length of I_k .

$$m(H)(1 - \Phi_{j_k}(H)b_k(x)) = \int_{|y-x|>r_k} m(H)(1 - \Phi_{j_k}(H)(x, y)b_k(y))dy$$

Apply weighted condition Lemma 3.1 (c)

$$\begin{aligned} & |\{x \notin \cup I_k^* : |\sum_k m(H)(1 - \Phi_{j_k}(H)b_k(x))| > \alpha/4\}| \\ & \leq C(\alpha/4)^{-1} \int_{\mathbb{R}^n \setminus \cup I_k^*} |\sum_k m(H)(1 - \Phi_{j_k}(H))b_k(x)|dx \\ & \leq C\alpha^{-1} \int \sum_k |b_k(y)|dy \int_{|y-x|>r_k} |m(H)(1 - \Phi_{j_k}(H)(x, y))|dx \\ & \leq C \sup_{\lambda} \|\chi m(\lambda \xi^2)\|_{C_{loc}^s} \alpha^{-1} \int |b(y)|dy \\ & \leq C \sup_{\lambda} \|\chi m(\lambda \xi^2)\|_{C_{loc}^s} \alpha^{-1} \|f\|_1. \end{aligned}$$

where we note

$$\begin{aligned} & \int_{|x-y|>r_k} |m(H)(1 - \Phi_{j_k}(H))(x, y)|dx \\ & \leq \sum_{2^j > r_k^2} \int_{|x-y|>r_k} |m_j(H)(x, y)|dx \leq C \end{aligned}$$

because if $\Phi(x) + \sum_{j=1}^{\infty} \phi(2^{-j}x) = 1$ then for any $j_0 \in \mathbb{Z}$, $\Phi(2^{-j_0}x) = 1 - \sum_{j=j_0+1}^{\infty} \phi(2^{-j}x)$.

Lower energy

Since $m(H)$ is bounded on L^2 . The proof is complete if we can show

$$(6) \quad \int |\sum_k \Phi_{j_k}(H)b_k(x)|^2 dx \leq C\alpha \|f\|_1$$

To show this let $h \in L^2(\mathbb{R}^n)$, $2^{-j_k} \sim \ell(I_k)^2$. According to (b)

$$\begin{aligned}
& \langle \sum_k \Phi_{j_k}(H)b_k, h \rangle \\
&= \sum_k \int_x h(x) dx \int_{y \in I_k} K_{j_k}(x, y) b_k(y) dy \\
&\leq \sum_k |I_k|^{-1} \int_x |h(x)| dx \int_{z \in I_k} \int_u \rho_j(x - z - u) d\mu(u) dz \int_y |b_k(y)| dy \\
&\leq \sum_k \|b_k\|_1 |I_k|^{-1} \int_{z \in I_k} \int_u (Mh)(z + u) d\mu dz \\
&\quad (\text{bec } \rho_j = 2^{jn/2} (1 + 2^{j/2}(\cdot))^{-n-\epsilon} \text{ is } \sim \text{ an approximation to the id}) \\
&\leq C\alpha \int_z \sum_k \chi_{I_k}(z) (Mh) * d\mu(z) dz \\
&\leq C\alpha \left\| \sum_k \chi_{I_k} \right\|_2 \|Mh * d\mu\|_2 \\
&\leq C\alpha \left(\sum_k |I_k| \right)^{1/2} \|h\|_2 \\
&\leq C\alpha (\alpha^{-1} \|f\|_1)^{1/2} \|h\|_2 = C\alpha^{1/2} \|f\|_1^{1/2} \|h\|_2
\end{aligned}$$

which proves (11). We have used the fact that if $\rho_t = t^{-n} \rho(x/t)$ is any approximation kernel to the identity, so that $\rho \in L^1(\mathbb{R}^n)$ is positive and decreasing, then

$$\sup_{t>0} |\rho_t * f(x)| \leq Mf(x)$$

where M denotes the Hardy-Littlewood maximal function on $\mathbb{R}^n$. $\square$

Remark. Note that in analyzing the kernel of $\Phi_j(H)$, oscillatory integral, if $j < 0$ we can prove the rapid decay from the spectrum side; if $j > 0$ it only gives a decay of $|x - y|^{-n}$; we will have to work in the space side and use the (average) integral version of rapid decay.

Remark. From the proof above we see the the weighted L^2 inequality in (b) somehow plays the role of Hörmander condition in classical case [37, 3]

$$(7) \quad \int_{|x-y|>2|x-\bar{y}|} |K(x, y) - K(x, \bar{y})| dx \leq A$$

which requires the gradient estimate for $K_j(x, y)$.

Remark. A nontrivial vector-valued version of the proof of the L^p result yields the multiplier result on the homogenous spaces $\dot{F}(H)$ and $\dot{B}(H)$ [31, 47].

In [30] when identifying the inhomogeneous space $F_p^{0,2}(H) = L^p$ we used decay for the derivative of the kernel in high energy, namely, (b') For $t > 0, j \geq 0$,

$$\int_{|x-y| \geq 2t} |m_j(H)(x, y) - m_j(H)(x, \bar{y})| dx \leq C(2^{j/2}t)^{\frac{1}{2}}$$

(c')

$$\int_{|x-y| \geq 2t} \sum_{2^j \geq t^2}^{\infty} |m_j(H)(x, y)| dx \leq C.$$

which is valid only for high energy estimate because we only have available for $j \geq 0$ the weighted estimate

$$\|(x - y)\partial_y K_j(x, y)\|_2 \leq 2^{j/4}.$$

To deal with the problem in low energy we avoid using the estimate for $\partial_y K_j(x, y)$ and follow the line of proof of Theorem 2.1 [31] for the homogeneous spaces $\dot{F}(H)$. Thus we obtain the identification of $\dot{F}(H)$ and L^p spaces.

Corollary 3.3. *Let $1 < p < \infty$. Then*

$$\dot{F}_p^{0,2}(H) = \dot{F}_p^{0,2}(\mathbb{R}^n) = L^p(\mathbb{R}^n),$$

meaning that the Littlewood-Paley characterization holds

$$\left\| \left(\sum_{j \in \mathbb{Z}} |\phi_j(H)f|^2 \right)^{1/2} \right\|_p \approx \|f\|_p.$$

4. PROOF OF COROLLARY 3.3

Identification of $\dot{F}_p^{\alpha,q}(H) = L^p$. *homogeneous spaces*

Let $Q_j = \phi_j(H)$, $j \in \mathbb{Z}$. Define

$$Q : f \mapsto \{\phi_j(H)f\}$$

and

$$R : \{f_j\} \mapsto \sum_{j=-\infty}^{\infty} \psi_j(H)f_j$$

We show that the same method in showing spectral multiplier theorem for F spaces yields

$$Q : L^1 \rightarrow w - L^1(\ell^2)$$

and

$$R : L^1(\ell^2) \rightarrow w - L^1$$

Hence, this, together with the boundedness $Q: L^2 \rightarrow L^2(\ell^2)$ and $R: L^2(\ell^2) \rightarrow L^2$, proves that if $1 < p < \infty$

$$\|f\|_{L^p} \sim \|\phi_j(H)f\|_{L^p(\ell^2)} = \|f\|_{\dot{F}_p^{0,2}(H)}$$

Lemma 4.1. $Q: L^2(\mathbb{R}^n) \rightarrow L^2(\ell^2)$. $R: L^2(\ell^2) \rightarrow L^2(\mathbb{R}^n)$.

Proof.

$$\begin{aligned} & \sum_{j \in \mathbb{Z}} (\phi_j(H)f, \phi_j(H)f) \\ &= \left(\sum_{j \in \mathbb{Z}} (\overline{\phi_j} \phi_j)(H)f, f \right) \lesssim \|f\|_2^2 \end{aligned}$$

because $\sum_{j \in \mathbb{Z}} |\phi_j(x)|^2 \approx 1$. The proof for R is similar. $\square$

Lemma 4.2. Q is $L^1 \rightarrow w - L^1(\ell^2)$, i.e.,

$$|\{ \sum_{j \in \mathbb{Z}} |\phi_j(H)f(x)|^2 \}^{1/2} > \alpha \}| \leq C\alpha^{-1} \|f\|_1$$

4.3. Proof of Q : weak $(L^1, L^1(\ell^2))$. Let $f \in L^1(\mathbb{R}^n)$. Given $\alpha > 0$, let $f = g + b$ be the Calderón-Zygmund decomposition, where $g \in L^2 \cap L^1$, $b = \sum_k b_k$, $b_k \subset I_k$, I_k disjoint.

- i) $|g(x)| \leq C\alpha$
- ii) $|I_k|^{-1} \int_{I_k} |f| dx \leq C_n \alpha$.

Note that (i), (ii) imply

$$|I_k|^{-1} \int_{I_k} |b| dx \leq C_n \alpha.$$

Since $\phi_j(H)$ is bounded $L^2 \rightarrow L^2(\ell^2)$.

$$\int \sum_{j \in \mathbb{Z}} |\phi_j(H)g(x)|^2 dx \leq C \|g\|_2^2 \leq C\alpha \|f\|_1$$

(i) gives

$$|\{x : (\sum_{j \in \mathbb{Z}} |\phi_j(H)g(x)|^2)^{1/2} > \alpha/2\}| \leq C \|\phi\|_\infty \alpha^{-1} \|f\|_1.$$

We estimate b in more detail. Let $2^{-j_k} \sim t_k = \text{diameter of the cube } I_k$.

$$\begin{aligned}
& |\{x : (\sum_{j \in \mathbb{Z}} |\phi_j(H)b(x)|^2)^{1/2} > \alpha/2\}| \\
& \leq |\{x : (\sum_j |\sum_k \phi_j(H)(1 - \Phi_{j_k}(H)b_k(x)|^2)^{1/2} > \alpha/4\}| \\
& + |\{x : (\sum_j |\sum_k \phi_j(H)\Phi_{j_k}(H)b_k(x)|^2)^{1/2} > \alpha/4\}| \\
& := B_1 + B_2.
\end{aligned}$$

For the first term since $|\cup I_k| \leq \alpha^{-1}\|f\|_1$, it suffices to estimate:

Denote I_k^* the cube having length three times the length of I_k with the same center as I_k . If $x \notin \cup_k I_k^*$, $I_k \subset \{y : |y - x| > r_k\}$, r_k being the length of I_k .

$$\phi_j(H)(1 - \Phi_{j_k}(H)b_k(x)) = \int_{|y-x|>r_k} \phi_j(H)(1 - \Phi_{j_k}(H)(x, y)b_k(y))dy$$

Apply Hebisch-Zheng condition (b)

$$\begin{aligned}
& |\{x \notin \cup_k I_k^* : (\sum_j |\sum_k \phi_j(H)(1 - \Phi_{j_k}(H)b_k(x)|^2)^{1/2} > \alpha/4\}| \\
& \leq C(\alpha/4)^{-1} \int_{\mathbb{R}^n \setminus \cup_k I_k^*} (\sum_j |\sum_k \phi_j(H)(1 - \Phi_{j_k}(H)b_k(x)|^2)^{1/2} dx \\
& \leq C\alpha^{-1} \int_{\mathbb{R}^n \setminus \cup_k I_k^*} \sum_k (\sum_j |\phi_j(H)(1 - \Phi_{j_k}(H)b_k(x)|^2)^{1/2} dx \quad (Minkowski) \\
& \leq C\alpha^{-1} \sum_k \int |b_k(y)|dy \int_{\mathbb{R}^n \setminus \cup_k I_k^*} \sum_{j \in \mathbb{Z}} |\phi_j(H)(1 - \Phi_{j_k}(H)(x, y))| dx \quad (Jensen) \\
& \leq C\alpha^{-1} \sum_k \int |b_k(y)|dy \int_{\mathbb{R}^n \setminus I_k} \sum_{j \geq j_k} |\phi_j(H)(1 - \Phi_{j_k}(H)(x, y))| dx \\
& \leq C \sup_{\lambda} \|\phi_j(\lambda \xi^2)\|_{H_{loc}^s} \alpha^{-1} \|f\|_1.
\end{aligned}$$

It remains to deal with B_2 . (Low energy) The proof is finished if we can show

$$(8) \quad \int \sum_j |\phi_j(H) \sum_k \Phi_{j_k}(H)b_k(x)|^2 dx \leq C\alpha \|f\|_1$$

Since $Q = \{\phi_j(H)\}_{j \in \mathbb{Z}}$ is bounded $L^2 \rightarrow L^2(\ell^2)$, we only need to prove

$$(9) \quad \int \sum_k |\Phi_{j_k}(H)b_k(x)|^2 dx \leq C\alpha \|f\|_1.$$

Similar to the proof of Theorem 2.1 we use duality argument. Let $h \in L^2(\mathbb{R}^n)$, $\|h\|_{L^2} \leq 1$. Then by condition (b)

$$\begin{aligned} & \langle \sum_k \Phi_{j_k}(H)b_k(x), h \rangle \\ & \leq c \sum_k \|b_k\|_1 |I_k|^{-1} \int_{z \in I_k} dz \int_u d\mu(u) \int_x \rho_t(x - z - u) |h(x)| dx \\ & = c \sum_k |I_k|^{-1} \|b_k\|_1 \int_z \chi_{I_k}(z) dz \int_u d\mu(u) \int_x \rho_t(x - z - u) |h(x)| dx \\ & \leq c\alpha \int_z \sum_k \chi_{I_k}(z) dz \int Mh(z + u) d\mu(u) \\ & \leq c\alpha \int_z \left\| \sum_k \chi_{I_k} \right\|_2 \|(Mh) * \mu\|_2 \\ & \leq c\alpha (\alpha^{-1} \|f\|_1^2)^{1/2} \|h\|_2 \leq c(\alpha \|f\|_1)^{1/2} \end{aligned}$$

which proves (9) hence (8). This completes the proof of Lemma 4.2. $\square$

Similarly we can show

Lemma 4.4. R is $L^1(\ell^2) \rightarrow w - L^1$, i.e.,

$$|\{x : |\sum_{j \in \mathbb{Z}} \psi_j(H)f_j(x)| > \alpha\}| \leq C\alpha^{-1} \|(\sum_j |f_j|^2)^{1/2}\|_1$$

Proof. Let $\{f_j\} \in L^1(\ell^2)$, $\alpha > 0$. Let $F(x) = (\sum_{j=-\infty}^{\infty} |f_j(x)|^2)^{1/2}$. By the C-Z decomposition for $F \in L^1$ there exists a collection of disjoint open cubes $\{I_k\}$ such that

$$\begin{aligned} i) \quad & |F(x)| \leq \alpha, \quad \text{a.e. } x \in \mathbb{R}^n \setminus \cup_k I_k \\ ii) \quad & \lambda \leq |I_k|^{-1} \int_{I_k} |F(x)| dx \leq 2^n \alpha, \quad \forall k. \end{aligned}$$

Define

$$g_j(x) = \begin{cases} |I_k|^{-1} \int_{I_k} f_j dx & x \in I_k \\ f_j(x) & \text{otherwise.} \end{cases}$$

and $b_j(x) = f_j(x) - g_j(x)$. Then $|g_j(x)| \leq C\alpha$.

Let $x \in I_k$. Minkowski's inequality gives

$$\begin{aligned} \left(\sum_{j=-\infty}^{\infty} |g_j(x)|^2 \right)^{1/2} &= \left(\sum_{j=-\infty}^{\infty} (|I_k|^{-1} \int_{I_k} |f_j| dx)^2 \right)^{1/2} \\ &\leq |I_k|^{-1} \int_{I_k} \left(\sum_{j=-\infty}^{\infty} |f_j|^2 \right)^{1/2} dx \leq 2^n \alpha. \end{aligned}$$

Thus $\int \sum_j |g_j|^2 \leq c\alpha \|F\|_1$ and by Lemma 4.1

$$\begin{aligned} |\{x : \sum_j |R_j g_j(x)|^2)^{1/2} > \alpha/2\}| &\leq C\alpha^{-2} \|\{R_j g_j\}\|_{L^2(\ell^2)}^2 \\ &\leq c\alpha^{-2} \|\{g_j\}\|_{L^2(\ell^2)}^2 \leq c\alpha^{-1} \|\{f_j\}\|_{L^1(\ell^2)}. \end{aligned}$$

(use the system $\{\phi_j\}_{-\infty}^{\infty}$)

It remains to estimate for $\{b_j(x)\}$. Let $2^{j_k} \sim \ell(I_k)^2$.

$$\begin{aligned} &\int_{\mathbb{R}^n \setminus I_k^*} \left(\sum_{j \in \mathbb{Z}} |R_j(1 - \Phi_{j_k}(H))b_{j,k}(x)|^2 \right)^{1/2} dx \\ &= \int_{\mathbb{R}^n \setminus I_k^*} \left(\sum_{j \sim j_k} \left| \int_{I_k} \psi_j(H)(1 - \Phi_{j_k}(H))(x, y) b_{j,k}(y) dy \right|^2 \right)^{1/2} dx \\ &\leq \int_{\mathbb{R}^n \setminus I_k^*} \int_{I_k} \left(\sum_j |\psi_j(H)(1 - \Phi_{j_k}(H))(x, y) b_{j,k}(y)|^2 \right)^{1/2} dy dx \quad \text{Minkowski} \\ &\leq \int_{\mathbb{R}^n \setminus I_k^*} \int_{I_k} \sup_j |\psi_j(H)(x, y)(1 - \Phi_{j_k}(H)(x, y))| \left(\sum_j |b_{j,k}(y)|^2 \right)^{1/2} dy dx \\ &\leq \int_{I_k} \left(\sum_j |b_{j,k}(y)|^2 \right)^{1/2} dy \int_{\mathbb{R}^n \setminus I_k^*} \sum_{j \geq j_k} |\psi_j(H)(1 - \Phi_{j_k}(H))(x, y)| dx \\ &\leq C \int_{I_k} \left(\sum_j |b_{j,k}(y)|^2 \right)^{1/2} dy \leq C' \int_{I_k} \left(\sum_j |f_j(y)|^2 \right)^{1/2} dy. \end{aligned}$$

where we used

$$\int_{|x-y| > \ell(I_k)} |\psi_j(H)(x, y)| dx \leq (2^{j/2} \ell(I_k))^{-1} \quad (N = n + 1)$$

by condition (1).

Hence,

$$\begin{aligned}
& |\{x \in \mathbb{R}^n \setminus \cup_k I_k^* : (\sum_j |R_j \sum_k (1 - \Phi_{j_k}(H)) b_{j,k}(x)|^2)^{1/2} > \alpha/2\}| \\
& \leq 2\alpha^{-1} \sum_k \int_{\mathbb{R}^n \setminus I_k^*} (\sum_j |R_j (1 - \Phi_{j_k}(H)) b_{j,k}(x)|^2)^{1/2} dx \\
& \leq C\alpha^{-1} \int (\sum_j |f_j(y)|^2)^{1/2} dy,
\end{aligned}$$

where $b_j = \sum_k b_{j,k}$ (convergence in $L^1 \cap L^2$ so $T_j b_j(x) = \sum_k T_j b_{j,k}$ in L^2) and we used

$$(\sum_j (\sum_k |T_j b_{j,k}(x)|^2)^{1/2})^{1/2} \leq \sum_k (\sum_j |T_j b_{j,k}(x)|^2)^{1/2}$$

by Minkowski inequality.

To estimate $|\{x \notin \cup_k I_k^* : (\sum_j |R_j \sum_k \Phi_{j_k}(H)) b_{j,k}(x)|^2)^{1/2} > \alpha/2\}|$, enough

$$(10) \quad \sum_j \int_{\mathbb{R}^n} |\psi_j(H) \sum_k \Phi_{j_k}(H) b_{j,k}(x)|^2 dx \leq C\alpha \|F\|_1.$$

Since $R = \{\psi_j(H)\}_{j \in \mathbb{Z}}$ is uniformly bounded $L^2 \rightarrow L^2$ or equivalently, $L^2(\ell^2) \rightarrow L^2(\ell^2)$, we only need show

$$(11) \quad \sum_j \int_{\mathbb{R}^n} |\sum_k \Phi_{j_k}(H) b_{j,k}(x)|^2 dx \leq C\alpha \|F\|_1.$$

Let $h = \{h_j\} \in L^2(\ell^2)$. $\|h\|_{L^2(\ell^2)} \leq 1$.

$$\begin{aligned}
& \sum_{j \in \mathbb{Z}} \langle \sum_k \Phi_{j_k}(H) b_{j,k}(x), h_j \rangle \\
& \leq C \sum_j \sum_k \|b_{j,k}\|_1 |I_k|^{-1} \int_{z \in I_k} dz \int_u d\mu(u) \int_x \rho_t(x - z - u) |h_j(x)| dx \\
& \leq C \sum_j \sum_k |I_k|^{-1} \|b_{j,k}\|_1 \int_z \chi_{I_k}(z) (\mu * Mh_j)(z) dz \\
& \leq C \left(\sum_j \left\| \sum_k |I_k|^{-1} \|b_{j,k}\|_1 \chi_{I_k} \right\|_2^2 \right)^{1/2} \|Mh_j\|_{L^2(\ell^2)} \\
& \leq C \left(\sum_j \sum_k |I_k|^{-1} \|b_{j,k}\|_1^2 \right)^{1/2} \|h_j\|_{L^2(\ell^2)} \\
& \leq C \left[\sum_k |I_k|^{-1} \sum_j \left(\int_{I_k} |b_j(y)| dy \right)^2 \right]^{1/2} \\
& \leq C \left[\sum_k |I_k|^{-1} \left(\int_{I_k} \left(\sum_j |b_j|^2 \right)^{1/2} dy \right)^2 \right]^{1/2} \leq C(\alpha \|F\|_1)^{1/2}.
\end{aligned}$$

□

where we have used the Fefferman-Stein inequality: if $1 < p < \infty$

$$\left\| \left(\sum_j (Mf_j)^2 \right)^{1/2} \right\|_p \leq \left\| \left(\sum_j |f_j|^2 \right)^{1/2} \right\|_p.$$

Remark. In the above proof of two lemmas we can replace $\{\phi_j\}_{j=-\infty}^{\infty}$ with $\{\Phi, \phi_j\}_{j=1}^{\infty}$ to obtain the inhomogeneous result. The homogeneous result is necessary and useful for obtaining Strichartz estimates for wave equation.

5. PÖSCHL-TELLER POTENTIAL

For $H = -d^2/dx^2 + V_\nu$ solve the Helmholtz equation

$$(12) \quad He(x, z) = z^2 e(x, z).$$

Under suitable asymptotic condition the solution also solves the Lippman-Schwinger equation

$$(13) \quad e(x, k) = e^{ikx} + \frac{1}{2ik} \int e^{ik|x-y|} V(y) e(y, k) dy,$$

that is

$$\begin{aligned} e(x, k) &= e^{ikx} - R_0(k^2 + i0)V e(\cdot, k)(x) \\ &= \sum_{n=0}^{\infty} (-R_0(k^2 + i0)V)^n e^{ikx} \\ &= (I + R_0(k^2 + i0)V)^{-1} e^{ikx}. \end{aligned}$$

Alternatively we can also using the above equations to write

$$e(x, k) = e^{ikx} - R_V(k^2)V e^{ikx}.$$

where the free resolvent has the kernel

$$R_0(k^2 + i0)(x, y) = -\frac{1}{2ik} e^{ik|x-y|} \quad k \in \mathbb{R} \setminus \{0\}.$$

From [30] we know that the continuous spectrum is $\sigma_c = [0, \infty)$, and the point spectrum is $\sigma_p = -1, -4, \dots, -n^2$. Bound states are Schwartz functions that are bounded by $c e^{-|x|}$.

We obtained in [30] the following formula for the continuum eigenfunctions.

Proposition 5.1. *Let $k \in \mathbb{R} \setminus \{0\}$. Then*

$$e_n(x, k) = (\text{sign}(k))^n \left(\prod_{j=1}^n \frac{1}{j + i|k|} \right) P_n(x, k) e^{ikx},$$

where $P_n(x, k) = p_n(\tanh x, ik)$ is defined by the recursion formula

$$p_n(\tanh x, ik) = \frac{d}{dx} (p_{n-1}(\tanh x, ik)) + (ik - n \tanh x) p_{n-1}(\tanh x, ik).$$

In particular, the function

$$\mathbb{R} \times (\mathbb{R} \setminus \{0\}) \ni (x, k) \mapsto e_n(x, k) \in \mathbb{C}$$

is analytic with $e_n(x, -k) = e_n(-x, k)$. Moreover, the function

$$(x, y, k) \mapsto e_n(x, k) \overline{e_n(y, k)} = \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k) P_n(y, -k) e^{ik(x-y)}$$

is real analytic on $\mathbb{R}^3$.

5.2. Resonance. The Wronskian can be computed by using Jost functions

$$W(z) = -2(-1)^n ik \prod_{\iota=1}^n \frac{\iota + ik}{\iota - ik}.$$

The remaining of this section is devoted to proving that conditions (a) and (b) in Theorem 2.2 are true for $H_\nu = H_0 + V_\nu$.

5.3. Weighted L^2 for $m(H)(x, y)$. Let $K_j(x, y)$ denote the kernel of $(m\phi_j)(H)E_{ac}$. Let $\lambda = 2^{-j/2}$.

Lemma 5.4.

$$(14) \quad \|(x - y)K_j(\cdot, y)\|_2 \leq \lambda^{1/2} \quad \forall j$$

$$(15) \quad \|(x - y)\partial_y K_j(\cdot, y)\|_2 \leq \lambda^{-1/2} \quad j \geq 0$$

Proof. Proof of (14).

$$\begin{aligned} & 2\pi i(x - y)K_j(\cdot, y) \\ &= i(x - y) \int m_j(k^2)e(x, k)\overline{e(y, k)}dk \\ &= \int_{|k| \sim \lambda^{-1}} m_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k)P_n(y, -k)\partial_k(e^{ik(x-y)})dk \\ &= - \int_{|k| \sim \lambda^{-1}} \partial_k[m_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k)P_n(y, -k)]e^{ik(x-y)}dk, \end{aligned}$$

which can be written as finite sums of

$$\begin{aligned} & (\tanh x)^\ell (\tanh y)^k [(m_j(k^2))' (\prod_{j=1}^n (j^2 + k^2))^{-1} r_{2n}(k)]^\vee (x - y) \\ & (\tanh x)^\ell (\tanh y)^k [m_j(k^2) (\prod_{j=1}^n (j^2 + k^2))^{-1} r_{2n-1}(k)]^\vee (x - y) \end{aligned}$$

and

$$(\tanh x)^\ell (\tanh y)^k [m_j(k^2) \frac{2k}{\iota^2 + k^2} (\prod_{j=1}^n (j^2 + k^2))^{-1} q_{2n}(k)]^\vee (x - y)$$

$0 \leq \ell, k, \iota \leq n$, $0 \leq i \leq 2n$, r_i, q_i are polynomials of degree i .

Plancherel formula for Fourier transform gives

$$\|(x - y)K_j(\cdot, y)\|_2 = O(\lambda^{1/2}) = O(2^{-j/4}) \quad \forall j.$$

using

$$\begin{cases} (m_j(k^2))^{(i)} = O(\frac{1}{k^i}) & i = 0, 1 \\ (\prod_{j=1}^n (j^2 + k^2))^{-1} r_i(k) = O(1/\langle k \rangle) \\ \frac{2k}{\iota^2 + k^2} = O(1/\langle k \rangle) \end{cases}$$

Proof of (15).

$$\begin{aligned}
& 2\pi i(x-y)\partial_y K_j(\cdot, y) \\
&= i(x-y) \int m_j(k^2) e(x, k) \overline{\partial_y e(y, k)} dk \\
&= \int_{|k| \sim \lambda^{-1}} (-ik) m_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k) P_n(y, -k) \partial_k (e^{ik(x-y)}) dk \\
&= \int_{|k| \sim \lambda^{-1}} m_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k) \partial_y (P_n(y, -k)) \partial_k (e^{ik(x-y)}) dk \\
&:= I_1 + I_2
\end{aligned}$$

$$I_1 = - \int_{|k| \sim \lambda^{-1}} \partial_k [-ik m_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k) P_n(y, -k)] e^{ik(x-y)} dk,$$

A similar argument as proving (14) yields

$$\|(x-y)K'_{1,j}(\cdot, y)\|_2 = O(\lambda^{-1/2}) = O(2^{j/4}) \quad \forall j.$$

using

$$\begin{cases} (m_j(k^2))^{(i)} = O(\frac{1}{k^i}) & i = 0, 1 \\ k = O(k) \\ (\prod_{j=1}^n (j^2 + k^2))^{-1} r_i(k) = O(1/\langle k \rangle) \\ \frac{2k}{i^2 + k^2} = O(1/\langle k \rangle) \end{cases}$$

$$\begin{aligned}
I_2 &= - \int_{|k| \sim \lambda^{-1}} \partial_k [m_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k) \partial_y (P_n(y, -k))] e^{ik(x-y)} dk \\
&= -\operatorname{sech}^2 y \int_{|k| \sim \lambda^{-1}} \partial_k [m_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k) (\partial_y p_n)(\tanh y, -k)] e^{ik(x-y)} dk.
\end{aligned}$$

where we find if $k \sim \lambda^{-1}$

$$\partial_k [m_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k) (\partial_y p_n)(\tanh y, -k)] = \begin{cases} O(k^{-2}) = O(1) & |k| \rightarrow \infty \\ O(k^{-1}) & |k| \rightarrow 0 \end{cases}$$

Plancherel formula for Fourier transform gives

$$\|(x-y)K'_{2,j}(\cdot, y)\|_2 = \operatorname{sech}^2 y \begin{cases} O(\lambda^{3/2}) = O(\lambda^{-1/2}) = O(2^{j/4}) & j \in \mathbb{N}_0 \\ O(\lambda^{1/2}) = O(2^{-j/4}) & j < 0 \end{cases}$$

□

Lemma 5.5. *Let $j \in \mathbb{Z}$.*

$$\begin{aligned} \|K_j(\cdot, y)\|_2 &\leq \lambda^{-1/2} = 2^{j/4} \quad \forall y \\ \|K_j(\cdot, y)\|_\infty &\lesssim 2^{j/2}. \end{aligned}$$

Proof. $K_j(x, y) = (m\phi_j)(H)E_{ac}(x, y)$.

$$\begin{aligned} 2\pi K_j(\cdot, y) &= \int m_j(k^2)e(x, k)\overline{e(y, k)}dk \\ &= \int_{|k|\sim\lambda^{-1}} m_j(k^2) \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) P_n(x, k)P_n(y, -k)(e^{ik(x-y)})dk \end{aligned}$$

which can be written as finite sums of

$$(\tanh x)^\ell (\tanh y)^k [(m_j(k^2))(\prod_{j=1}^n (j^2 + k^2))^{-1} r_{2n}(k)]^\vee (x - y)$$

$\therefore$

$$\begin{aligned} \|m_j(H)(x, y)\|_2 &\leq \|m_j(k^2)(\prod_{j=1}^n (j^2 + k^2))^{-1} r_{2n}(k)\|_{L_k^2} \\ &\sim \|m_j(k^2)\|_{L_k^2} \sim 2^{j/4} (\int_{|k|\leq 1} |\phi(k^2)|^2 dk)^{1/2}, \quad \forall j \end{aligned}$$

(if $m = 1$, $\phi_j \subset [-2^j, 2^j]$), using

$$\begin{cases} m_j(k^2) = O(1) \\ r_{2n}(k) \prod_{j=1}^n (j^2 + k^2)^{-1} = O(1) \end{cases}$$

□

Lemma 5.6.

$$\begin{aligned} \|(x - y)\partial_y K_j(\cdot, y)\|_2 &\lesssim \begin{cases} 2^{j/4} + \operatorname{sech}^2 y 2^{-j/4} & j \rightarrow -\infty \\ 2^{j/4} & j \rightarrow \infty \end{cases} \\ \|(x - y)\partial_y K_j(\cdot, y)\|_\infty &\lesssim 2^{j/2} + \operatorname{sech}^2 y \begin{cases} O(1) & j \rightarrow -\infty \\ 2^{-j/2} & j \rightarrow \infty \end{cases} \end{aligned}$$

Remark. This means for $j \rightarrow -\infty$, $\|(x - y)\partial_y K_j(\cdot, y)\|_r \sim \operatorname{sech}^2 y 2^{-j/(2r)}$, $r \in [2, \infty]$, which does not seem to help establish the Hörmander integral condition even if using r -norm instead of 2-norm.

Proof. For 2-norm, it is proved before. For $r = \infty$,

$$\begin{aligned}
& i2\pi(x-y)\partial_y K_j(\cdot, y) \\
&= - \int \partial_k [km_j(k^2)] \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) p_n(\tanh x, k) p_n(\tanh y, -k) e^{ik(x-y)} dk \\
&+ (-i) \operatorname{sech}^2 y \int_{|k| \sim \lambda^{-1}} \partial_k [m_j(k^2)] \left(\prod_{j=1}^n \frac{1}{j^2 + k^2} \right) p_n(\tanh x, k) (\partial_y p_n)(\tanh y, -k) e^{ik(x-y)} dk \\
&\sim \int_{|k| \sim \lambda^{-1}} O(1) dk + \operatorname{sech}^2 y \int_{|k| \sim \lambda^{-1}} \begin{cases} O(1/k) & k \rightarrow 0 \\ O(1/k^2) = O(1) & k \rightarrow \infty \end{cases} dk \\
&(\lambda = 2^{-j/2}) \\
&\therefore
\end{aligned}$$

$$\|(x-y)\partial_y K_j(\cdot, y)\|_\infty \sim 2^{j/2} + \operatorname{sech}^2 y \begin{cases} O(1) & j \rightarrow -\infty \\ 2^{-j/2} & j \rightarrow \infty. \end{cases}$$

□

5.7. Weighted pointwise decay of the kernel. The problem for pointwise decay of $\Phi_j(H)(x, y)$ in higher energy can be overcome by using an integral version of (3) with a finite measure¹.

Lemma 5.8. (*Hebisch-Zheng*) Let $\Psi \in C_0^\infty$ be supported in $[-1, 1]$ and let I be any cube in $\mathbb{R}^n$ with length $\ell(I)$. Then for all $x \in \mathbb{R}^n$ and $y \in I$ with $\ell(I) = 2^{-j/2}$ we have

a)

$$|\Phi_j(H)(x, y)| \leq c \int_{u \in \mathbb{R}^n} \frac{2^{jn/2}}{(1 + 2^{j/2}|x - y - u|)^{n+\epsilon}} d\mu(u)$$

$$d\mu(u) = \delta(u) + \langle u \rangle^m e^{-c|u|} du, \text{ some } m \geq 0.$$

Hence b)

$$\sup_{y \in I} |\Phi_j(H)(x, y)| \leq \frac{c}{|I|} \int_{z \in I} \int_{u \in \mathbb{R}^n} \frac{2^{jn/2}}{(1 + 2^{j/2}|x - z - u|)^{n+\epsilon}} d\mu(u) dz.$$

¹ We modify Hebisch method when the kernel is rough (and slowly decaying), not having Lipschitz smoothness as needed in the Hörmander method

Proof. Let $\Psi(x) = \Phi(x^2)$, $\lambda = 2^{-j/2}$. According to the formula for the kernel in preceding subsection

$$\begin{aligned}
& \Phi_j(H)(x, y) \\
&= [\Psi(\lambda k)]^\vee(x - y) + [\Psi(\lambda k) \sum_{\substack{\iota=1, \dots, n \\ \mu_\iota=0, \dots, N}} \frac{a_{\mu_\iota, \iota} + b_{\mu_\iota, \iota} k}{(\iota^2 + k^2)^{\mu_\iota+1}}]^\vee(x - y) \\
&= \lambda^{-1} \Psi^\vee(\lambda^{-1}(x - y)) \\
&+ c \lambda^{-1} \int_{\mathbb{R}} \Psi^\vee(\lambda^{-1}(x - y - u)) |u|^m e^{-c|u|} du,
\end{aligned}$$

$m \in \mathbb{N}_0$ is an integral constant. Thus

$$\begin{aligned}
& |\Phi_j(H)(x, y)| = \left| \int_{|k| \lesssim 2^{j/2}} \Psi(2^{-j/2} k) e^{i(x-y)k} (1 + \hat{a}(k)) dk \right| \\
&= (\lambda^{-n} \Psi^\vee(\lambda^{-1} \cdot) * (1 + \hat{a})^\vee)(x - y) \\
&\lesssim \lambda^{-n} \int_{\mathbb{R}^n} (1 + \lambda^{-1} |x - y - u|)^{-n-\epsilon} d\mu(u),
\end{aligned}$$

where $y \in I$, $d\mu(u) = \delta(u) + a(u)du$, $a(u) = |u|^m e^{-c|u|}$. $\square$

Remark. Observe that for $j > 0$, Φ_j actually contains both high+low energy information (if $0 \in \text{supp } \Phi$). This is the most technically difficult part. Fortunately with Lemma 5.8 we can control it by maximal function.

5.9. Kernel decay for a positive potential. The case is simpler when V is nonnegative. It can be shown [46, 9] that (3) is a weaker assumption than the heat kernel estimate

$$0 \leq e^{-tH}(x, y) \leq c_n t^{-n/2} e^{-cd(x,y)^2/t} \quad \forall t > 0.$$

Examples include H being a uniform elliptic operator or its perturbation of order 0. i.e., a Schrödinger operator with $V = V_+ - V_-$, V_- is small in Kato norm, cf. [8].

In [32] it is shown if $\sigma(H) \subset [0, \infty)$ then

$$e^{-tH}(x, y) \leq C t^{-n/2} e^{-|x-y|^2/4t} (1 + \delta t + |x-y|^2/t)^{n/2} \quad \forall t > 0$$

$\delta = \delta(V_-)$ and $\delta = 0$ if $V_- = 0$.

Proposition 5.10. *Let H denote a selfadjoint operator on (M, g) with dimension n . Suppose e^{-tH} verifies the upper Gaussian bound*

$$e^{-tH}(x, y) \leq C_n t^{-n/2} e^{-cd(x,y)^2/t} \quad \forall t > 0.$$

Then for each ℓ

$$|\phi_j(H)(x, y)| \leq C_\ell 2^{jn/2} (1 + 2^{j/2} d(x, y))^{-\ell} \quad \forall j \in \mathbb{Z}.$$

REFERENCES

- [1] S. Agmon, Spectral properties of Schrödinger operators and scattering theory, *Annali Scuola Norm. Sup. de Pisa* **2** (1975), 151–218.
- [2] G. Alexopoulos, Spectral multipliers on Lie groups of polynomial growth, *Proc. A.M.S.* **120** (1994) no.3, 973–979.
- [3] J.J. Benedetto, S. Zheng, Besov spaces for the Schrödinger operator with barrier potential (submitted). <http://lanl.arXiv.org/math.CA/0411348>, (2005).
- [4] J. Bergh, J. Löfström, *Interpolation Spaces*, Springer-Verlag, 1976.
- [5] H.R. Beyer, On the completeness of the quasinormal modes of the Pöschl-Teller potential, *Comm. Math. Phys.* **204** (1999), no. 2, 397–423.
- [6] F.M. Christ, C.D. Sogge, The weak type L^1 convergence of eigenfunction expansions for pseudodifferential operators, *Invent. Math.* **94** (2) (1988) 421–453.
- [7] M. Cowling, A. Sikora, A spectral multiplier theorem for a sublaplacian on $SU(2)$, *Math. Z.* **238** (2001), no. 1, 1–36.
- [8] P. D’Ancona, V. Pierfelice, On the wave equation with a large rough potential. *J. F. A.* <http://arXiv.org/math.AP/0310199>.
- [9] E.B. Davies, *Heat Kernels and Spectral Theory*, Cambridge University Press, Cambridge, 1989.
- [10] P. Deift, E. Trubowitz, Inverse scattering on the line. *Comm. Pure Appl. Math.* vol. XXXII. (1979). 121–251.
- [11] X.T. Duong, A. McIntosh, Singular integral operators with non-smooth kernels on irregular domains, *Rev. Mat. Iberoamericana*, **15** (1999), no.2: 233–265.
- [12] X.T. Duong, E.M. Ouhabaz and A. Sikora, Plancherel type estimates and sharp spectral multipliers, *J. Funct. Anal.* (2002).
- [13] X. T. Duong, L. Yan, Duality of Hardy and BMO spaces associated with operators with heat kernel bounds, *J. Amer. Math. Soc.* **18** (2005), 943–973.
- [14] J. Dziubański, Spectral multiplier theorem for H^1 spaces associated with some Schrödinger operators, *Proc. A.M.S.* **127** (1999), no. 12, 3605–3613.
- [15] ———, A spectral multiplier theorem for H^1 spaces associated with Schrödinger operators with potentials satisfying a reverse Hölder inequality. *Illinois J. Math.* **45** (2001), no. 4, 1301–1313.
- [16] J. Epperson, Hermite multipliers and pseudo-multipliers, *Proc. Amer. Math. Soc.* **124** (1996), no. 7, 2061–2068.
- [17] —, Hermite and Laguerre wave packet expansions. *Studia Math.* **126** (1997), no. 3, 199–217.
- [18] C. Fefferman, Inequalities for strongly singular convolution operators, *Acta Math.* **124** (1970), 9–36.
- [19] S. Flügge, *Practical Quantum Mechanics*, Springer-Verlag, 1974.
- [20] M. Frazier, B. Jawerth, G. Weiss, *Littlewood-Paley Theory and the Study of Function Spaces*, Conference Board of the Math. Sci. **79**, 1991.
- [21] C.-A. Guerin, M. Holschneider, Time-dependent scattering on fractal measures, *J. Math. Physics* **39**(8), 1998.
- [22] W. Hebisch: A multiplier theorem for Schrödinger operators. *Colloq. Math.* **60/61** (1990), no. 2, 659–664.

- [23] ———, Almost everywhere summability of eigenfunction expansions associated to elliptic operators. *Studia Math* **96** (1990), no. 3, 263–275.
- [24] ———, Functional calculus for slowly decaying kernels, preprint, 1995.
<http://www.math.uni.wroc.pl/~hebisch>
- [25] A. Jensen, S. Nakamura, Mapping properties of functions of Schrödinger operators between L^p spaces and Besov spaces, in *Spectral and Scattering Theory and Applications*, Advanced Studies in Pure Math. **23**, 1994.
- [26] S. Klainerman, M. Machedon and J. Stalker, Decay of solutions to the wave equation on a spherically symmetric static background, *preprint*.
- [27] G.L. Lamb, Jr., *Elements of Soliton Theory*, Pure & Applied Mathematics, Wiley-Interscience, 1980.
- [28] S.G. Mikhlin, *Multidimensional Singular Integrals and Integral Equations*, Pergamon Press, Oxford, 1965.
- [29] D. Müller, E.M. Stein, On spectral multipliers for Heisenberg and related groups, *J. Math. Pures Appl.* (9) **73** (4) (1994) 413–440.
- [30] G. Ólafsson, S. Zheng, Function spaces associated with Schrödinger operators: the Pöschl-Teller potential. accepted for publication in *Journal of Fourier Analysis and Applications*.
- [31] G. Ólafsson, K. Osolkov, S. Zheng, Spectral multipliers for Schrödinger operators: II, Preprint.
- [32] E.M. Ouhabaz, Sharp Gaussian bounds and L^p -growth of semigroups associated with elliptic and Schrödinger operators. *Proc. A.M.S.* 2006.
- [33] ———, *Analysis of Heat Equations on Domains*, London Math. Soc. Monographs, Vol. 31. Princeton Univ. Press 2004.
- [34] W. Schlag, Dispersive estimates for Schrödinger operators: A survey. <http://lanl.arXiv.org/math.AP/0501037>, (2005).
- [35] ———, A remark on Littlewood-Paley theory for the distorted Fourier transform, <http://lanl.arXiv.org/math.AP/0508577>, (2005)
- [36] B. Simon, Schrödinger semigroups, *Bull. Amer. Math. Soc.* **7** (1982) no.3, 447–526.
- [37] E. Stein, *Harmonic analysis, Real-variable methods, orthogonality, and oscillatory integrals*, Princeton Univ. Press, 1993.
- [38] M. Taylor, L^p -estimates on functions of the Laplace operator, *Duke Math. J.* **58** no. 3 (1989), 773–793
- [39] H. Triebel, *Theory of Function Spaces*, Birkhäuser Verlag, 1983.
- [40] —, *Theory of Function Spaces II*, Monographs Math. **84**, Birkhäuser, Basel, 1992.
- [41] R. Weder, The $W^{k,p}$ -continuity of the Schrödinger wave operators on the line. *Comm. Math. Phys.* **208** (1999) 507–520.
- [42] Yajima, K., The $W^{k,p}$ -continuity of wave operators for Schrödinger operators. *J. Math. Soc. Japan* **47**, 551–581 (1995).
- [43] ———, Dispersive estimate for Schrödinger equations with threshold resonance and eigenvalue. *Comm. Math. Phys.* 2005.
- [44] Q. Zhang, Global bounds of Schrödinger heat kernels with negative potentials. *J. Func. Anal.* **182** (2001), no.2, 344–370.
- [45] S. Zheng, A representation formula related to Schrödinger operators. *Anal. Theo. Appl.* **20** (2004), no.3. <http://lanl.arXiv.org/math.SP/0412314>.
- [46] ———, Littlewood-Paley theorem for Schrödinger operators. (sumitted) 2006.

- [47] ———, Spectral multipliers, function spaces and dispersive estimates for Schrödinger operators. *Preprint*.

DEPARTMENT OF MATHEMATICS, INDUSTRIAL MATHEMATICS INSTITUTE, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SC 29208

E-mail address: `shijun@math.sc.edu`

URL: `http://www.math.sc.edu/~shijun`